

Development

Volume 111 1991

Editor in Chief

Chris Wylie (Cambridge)

European Editors

Richard Gardner (Oxford)
Walter Gehring (Basel)

Peter Lawrence (Cambridge)
Keith Roberts (Norwich)

USA Editors

Doug Melton (Harvard)
Tom Jessell (Columbia)

Gerry Rubin (Berkeley)

Editorial Advisory Board

Eileen Adamson (La Jolla)
Michael Akam (Cambridge)
Michael Bate (Cambridge)
Rosa Beddington (Edinburgh)
Marianne Bronner-Fraser (Irvine)
Clayton Buck (Wistar Institute)
Alan Colman (Birmingham)
Eric Davidson (Pasadena)
Alun Davies (London)
Françoise Dieterlen (Nogent-sur-Marne)
John Ellis (Warwick)
Gary Felsenfeld (Bethesda)
Richard Firtel (San Diego)
Joseph Frankel (Iowa)
Michael Freeling (Berkeley)
John Gerhart (Berkeley)
Peter Goodfellow (London)
Peter Gruss (Göttingen)
John Gurdon (Cambridge)
John Heath (Oxford)
Brigid Hogan (Vanderbilt)
Nigel Holder (London)
Robert Horvitz (MIT)
Phil Ingham (Oxford)
Herbert Jäckle (Munich)
Rudolph Jaenisch (Whitehead, MIT)
William Jeffery (Austin)
Martin Johnson (Cambridge)
Klaus Kalthoff (Austin)

Robert Kay (Cambridge)
Judith Kimble (Madison)
Charles Kimmel (Eugene)
Klaus Kratochwil (Salzburg)
Ruth Lehmann (Whitehead, MIT)
Julian Lewis (Oxford)
Andrew Lumsden (London)
Mary Lyon (Didcot)
Terry Magnuson (Case Western)
Gail Martin (San Francisco)
Bjorn Olsen (Harvard)
Scott Poethig (Pennsylvania)
Martin Raff (London)
Elizabeth Robertson (Columbia)
Janet Rossant (Toronto)
Joan Ruderman (Duke)
Matthew Scott (Stanford)
Jonathan Slack (Oxford)
Jim Smith (London)
Michael Sporn (Bethesda)
Claudio Stern (Oxford)
Gary Struhl (Columbia)
Chris Summerville (Michigan)
Azim Surani (Cambridge)
Masatoshi Takeichi (Kyoto)
Peter Thorogood (London)
David Weisblat (Berkeley)
Richard Whittaker (Woods Hole)
Jeff Williams (ICRF, Clare Hall)

The Company of Biologists Limited, founded in 1925, is a 'Company Limited by Guarantee' having tax-exempt charitable status. There is a Board of Directors consisting of about 20 professional biologists, two of them appointed annually by the Society for Experimental Biology, who receive no salary or fees for their services. The Company's main function is to own and produce *The Journal of Experimental Biology*, the *Journal of Cell Science* and *Development*, and to appoint the Editors of these journals. These are part-time appointments held by established professional biologists of some eminence, and once they have been appointed the Company exercises no control over editorial policy.

The Company is precluded by its charitable status from making a commercial profit on its operations, and its aim is to produce high-quality journals at the lowest possible price. Any surplus on publishing not required for the journals' reserves is transferred to an Educational Trust Fund, which makes substantial grants in aid of societies concerned with the fields of interest covered by the Company's journals. Grants are also made to conferences and summer schools in the fields of its journals.

Typeset, Printed and Published by
The Company of Biologists Limited
Department of Zoology, University of Cambridge, Downing Street,
Cambridge CB2 3EJ

© The Company of Biologists Limited 1991

Contents

Volume 111 1991

Young, P. E., Pesacreta, T. C. and Kiehart, D. P. Dynamic changes in the distribution of cytoplasmic myosin during <i>Drosophila</i> embryogenesis	1	Cunha, G. R., Young, P., Higgins, S. J. and Cooke, P. S. Neonatal seminal vesicle mesenchyme induces a new morphological and functional phenotype in the epithelia of adult ureter and ductus deferens	145
Ranscht, B. and Bronner-Fraser, M. T-cadherin expression alternates with migrating neural crest cells in the trunk of the avian embryo	15	Levi, G., Gumbiner, B. and Thiery, J. P. The distribution of E-cadherin during <i>Xenopus laevis</i> development	159
Condic, M. L., Fristrom, D. and Fristrom, J. W. Apical cell shape changes during <i>Drosophila</i> imaginal leg disc elongation: a novel morphogenetic mechanism	23	Sun, F. Z. and Moor, R. M. Nuclear-cytoplasmic interactions during ovine oocyte maturation	171
Maden, M., Hunt, P., Eriksson, U., Kuroiwa, A., Krumlauf, R. and Summerbell, D. Retinoic acid-binding protein, rhombomeres and the neural crest	35	de Pomerai, D. I., Ip, W. K., McLaughlin, M. and Perry, K. C. Expression in non-lens tissues of an enzyme activity related to the 'lens-specific' protein, δ crystallin	181
Ruberte, E., Dollé, P., Chambon, P. and Morriss-Kay, G. Retinoic acid receptors and cellular retinoid binding proteins. II. Their differential pattern of transcription during early morphogenesis in mouse embryos	45	Linsenmayer, T. F., Chen, Q., Gibney, E., Gordon, M. K., Marchant, J. K., Mayne, R. and Schmid, T. M. Collagen types IX and X in the developing chick tibiotarsus: analyses of mRNAs and proteins	191
Murphy, P. and Hill, R. E. Expression of the mouse <i>labial</i> -like homeobox-containing genes, <i>Hox 2.9</i> and <i>Hox 1.6</i> , during segmentation of the hindbrain	61	Cooke, J. and Wong, A. Growth-factor-related proteins that are inducers in early amphibian development may mediate similar steps in amniote (bird) embryogenesis	197
Sobis, H., Verstuyf, A. and Vandeputte, M. Endodermal origin of yolk-sac-derived teratomas	75	Trautman, M. S., Kimelman, J. and Bernfield, M. Developmental expression of syndecan, an integral membrane proteoglycan, correlates with cell differentiation	213
Prokop, A. and Technau, G. M. The origin of postembryonic neuroblasts in the ventral nerve cord of <i>Drosophila melanogaster</i>	79	Bloch-Gallego, E., Huchet, M., El M'Hamdi, H., Xie, F.-K., Tanaka, H. and Henderson, C. E. Survival <i>in vitro</i> of motoneurons identified or purified by novel antibody-based methods is selectively enhanced by muscle-derived factors	221
Bang, A. G., Hartenstein, V. and Posakony, J. W. <i>Hairless</i> is required for the development of adult sensory organ precursor cells in <i>Drosophila</i>	89	Lyons, G. E., Buckingham, M. E., Tweedie, S. and Edwards, Y. H. Carbonic anhydrase III, an early mesodermal marker, is expressed in embryonic mouse skeletal muscle and notochord	233
Ayer-Le Lievre, C., Ståhlbom, P.-A. and Sara, V. R. Expression of IGF-I and -II mRNA in the brain and craniofacial region of the rat fetus	105	Smith, S. M. and Eichele, G. Temporal and regional differences in the expression pattern of distinct retinoic acid receptor- β transcripts in the chick embryo	245
Schmid, P., Cox, D., Bilbe, G., Maier, R. and McMaster, G. K. Differential expression of TGF $\beta 1$, $\beta 2$ and $\beta 3$ genes during mouse embryogenesis	117	Fisher, C., Byers, M. R., Iadarola, M. J. and Powers, E. A. Patterns of epithelial expression of Fos protein suggest important role in the transition from viable to cornified cell during keratinization	253
Millan, F. A., Denhez, F., Kondaiah, P. and Akhurst, R. J. Embryonic gene expression patterns of TGF $\beta 1$, $\beta 2$ and $\beta 3$ suggest different developmental functions <i>in vivo</i>	131		

Contents

Wiles, M. V. and Keller G. Multiple hematopoietic lineages develop from embryonic stem (ES) cells in culture	259	Sánchez-Herrero, E. Control of the expression of the bithorax complex genes abdominal-A and Abdominal-B by <i>cis</i> -regulatory regions in <i>Drosophila</i> embryos	437
Mackenzie, A., Leeming, G. L., Jowett, A. K., Ferguson, M. W. J. and Sharpe, P. T. The homeobox gene Hox 7.1 has specific regional and temporal expression patterns during early murine craniofacial embryogenesis, especially tooth development <i>in vivo</i> and <i>in vitro</i>	269	Lyons, G. E., Buckingham, M. E. and Mannherz, H. G. α -actin proteins and gene transcripts are colocalized in embryonic mouse muscle	451
Davis, C. A., Holmyard, D. P., Millen, K. J. and Joyner, A. L. Examining pattern formation in mouse, chicken and frog embryos with an <i>En</i> -specific antiserum	287	Wanaka, A., Milbrandt, J. and Johnson, E. M. Expression of FGF receptor gene in rat development	455
Fernandez, D. E., Turner, F. R. and Crouch, M. L. <i>In situ</i> localization of storage protein mRNAs in developing meristems of <i>Brassica napus</i> embryos	299	Drysdale, T. A. and Elinson, R. P. Development of the <i>Xenopus laevis</i> hatching gland and its relationship to surface ectoderm patterning	469
Ginsberg, D., DeSimone, D. and Geiger, B. Expression of a novel cadherin (EP-cadherin) in unfertilized eggs and early <i>Xenopus</i> embryos	315	Perkins, A. S., Mercer, J. A., Jenkins, N. A. and Copeland, N. G. Patterns of <i>Evi-1</i> expression in embryonic and adult tissues suggest that <i>Evi-1</i> plays an important regulatory role in the mouse development	479
Takahashi, Y. and Nogawa, H. Branching morphogenesis of mouse salivary epithelium in basement membrane-like substratum separated from mesenchyme by the membrane filter	327	Brown, R. and Brockes, J. P. Identification and expression of a regeneration-specific homeobox gene in the newt limb blastema	489
Shi, X., Lu, L., Qiu, Z., He, W. and Frankel, J. Microsurgically generated discontinuities provoke heritable changes in cellular handedness of a ciliate, <i>Stylonychia mytilus</i>	337	Ferretti, P., Brockes, J. P. and Brown, R. A new type II keratin restricted to normal and regeneration limbs and tails is responsive to retinoic acid	497
Boulter, C. A., Aguzzi, A., Williams, R. L., Wagner, E. F., Evans, M. J. and Beddington, R. Expression of <i>v-src</i> induces aberrant development and twinning in chimaeric mice	357	Minkoff, R., Parker, S. B. and Hertzberg, E. L. Analysis of distribution patterns of gap junctions during development of embryonic chick facial primordia and brain	509
Eldon, E. D. and Pirrotta, V. Interactions of the <i>Drosophila</i> gap gene <i>giant</i> with maternal and zygotic pattern-forming genes	367	Godsave, S. F. and Slack, J. M. W. Single cell analysis of mesoderm formation in the <i>Xenopus</i> embryo	523
Yasuda, G. K., Baker, J. and Schubiger, G. Independent roles of centrosomes and DNA in organizing the <i>Drosophila</i> cytoskeleton	379	Jones, C. M., Lyons, K. M. and Hogan, B. L. M. Involvement of <i>Bone Morphogenetic Protein-4</i> (BMP-4) and <i>Vgr-1</i> in morphogenesis and neurogenesis in the mouse	531
Boulet, A. M., Lloyd, A. and Sakonju, S. Molecular definition of the morphogenetic and regulatory functions and the <i>cis</i> -regulatory elements of the <i>Drosophila</i> <i>Abd-B</i> homeotic gene	393	Miwa, Y., Atsumi, T., Imai, N. and Ikawa, Y. Primitive erythropoiesis of mouse teratocarcinoma stem cells PCC3/A/1 in serum-free medium	543
Irvine, K. D., Helfand, S. L. and Hogness, D. S. The large upstream control region of the <i>Drosophila</i> homeotic gene <i>Ultrabithorax</i>	407	Hopwood, N. D., Pluck, A. and Gurdon, J. B. <i>Xenopus</i> Myf-5 marks early muscle cells and can activate muscle genes ectopically in early embryos	551
Swalla, B. J., Badgett, M. R. and Jeffery, W. R. Identification of a cytoskeletal protein localized in the myoplasm of ascidian eggs: localization is modified during anural development	425	Mazarakis, N. D., Nelki, D., Lyon, M. F., Ruddy, S., Evans, E. P., Freemont, P. and Dudley, K. Isolation and characterisation of a testis-expressed developmentally regulated gene from the distal inversion of the mouse <i>t</i> -complex	561

Sasaki, H., Hamada, T., Ueda, T., Seki, R., Higashinakagawa, T. and Sakaki, Y. Inherited type of allelic methylation variations in a mouse chromosome region where an integrated transgene shows methylation imprinting	573	Desbiens, X., Quéva, C., Jaffredo, T., Stéhelin, D. and Vandenbunder, B. The relationship between cell proliferation and the transcription of the nuclear oncogenes <i>c-myc</i> , <i>c-myb</i> and <i>c-ets-1</i> during feather morphogenesis in the chick embryo	699
Perris, R., Krotoski, D., Lallier, T., Domingo, C., Sorrell, J. M. and Bronner-Fraser, M. Spatial and temporal changes in the distribution of proteoglycans during avian neural crest development	583	Hemmati-Brivanlou, A., de la Torre, J. R., Holt, C. and Harland, R. M. Cephalic expression and molecular characterization of <i>Xenopus En-2</i>	715
Kraut, R. and Levine, M. Spatial regulation of the gap gene <i>giant</i> during <i>Drosophila</i> development	601	Hornbruch, A. and Wolpert, L. The spatial and temporal distribution of polarizing activity in the flank of the pre-limb-bud stages in the chick embryo	725
Kraut, R. and Levine, M. Mutually repressive interactions between the gap genes <i>giant</i> and <i>Krüppel</i> define middle body regions of the <i>Drosophila</i> embryo	611	Dudley, M. and Poethig, R. S. The effect of a heterochronic mutation, <i>Teopod2</i> , on the cell lineage of the maize shoot	733
Kelso-Winemiller, L. C. and Winkler, M. M. 'Unmasking' of stored maternal mRNAs and the activation of protein synthesis at fertilization in sea urchins	623	Moore, J. W., Dionne, C., Jaye, M. and Swain, J. L. The mRNAs encoding acidic FGF, basic FGF, and FGF receptor are coordinately downregulated during myogenic differentiation	741
Stocker, K. M., Sherman, L., Rees, S. and Ciment, G. Basic FGF and TGF- β 1 influence commitment to melanogenesis in neural-crest-derived cells of avian embryos	635	Thomas, U., Speicher, S. A. and Knust, E. The <i>Drosophila</i> gene <i>Serrate</i> encodes an EGF-like transmembrane protein with a complex expression pattern in embryos and wing discs	749
Pomeranz, H. D., Rothman, T. P. and Gershon, M. D. Colonization of the post-umbilical bowel by cells derived from the sacral neural crest: direct tracing of cell migration using an intercalating probe and a replication-deficient retrovirus	647	Kubiak, J., Paldi, A., Weber, M. and Maro, B. Genetically identical parthenogenetic mouse embryos produced by inhibition of the first meiotic cleavage with cytochalasin D	763
Blackman, R. K., Sanicola, M., Raftery, L. A., Gillevet, T. and Gelbart, W. M. An extensive 3' cis-regulatory region directs the imaginal disk expression of <i>decapentaplegic</i> , a member of the TGF- β family in <i>Drosophila</i>	657	Rowe, A., Eager, N. S. C. and Brickell, P. M. A member of the RXR nuclear receptor family is expressed in neural-crest-derived cells of the developing chick peripheral nervous system	771
Goh, P.-Y. and Bogaert, T. Positioning and maintenance of embryonic body wall muscle attachments in <i>C. elegans</i> requires the <i>mup-1</i> gene	667	Jermyn, K. A. and Williams, J. G. An analysis of culmination in <i>Dictyostelium</i> using prestalk and stalk-specific cell autonomous markers	779
Flamme, I., Schulze-Osthoff, K. and Jacob, H. J. Mitogenic activity of chicken chorioallantoic fluid is temporally correlated to vascular growth in the chorioallantoic membrane and related to fibroblast growth factors	683	Schlosshauer, B., Grauer, D., Dütting, D. and Vanselow, J. Expression of a novel Müller glia specific antigen during development and after optic nerve lesion	789
Andujar, M. B., Couble, P., Couble, M.-L. and Magloire, H. Differential expression of type I and type III collagen genes during tooth development	691	Radice, G., Lee, J. J. and Constantini, F. H β 58, an insertional mutation affecting early postimplantation development of the mouse embryo	801
		Jessus, C., Rime, H., Haccard, O., van Lint, J., Goris, J., Merlevede, W. and Ozon, R. Tyrosine phosphorylation of p34 ^{cdc2} and p42 during meiotic maturation of <i>Xenopus</i> oocyte. Antagonistic action of okadaic acid and 6-DMAP	813

Contents

Saiga, H., Mizokami, A., Makabe, K. W., Satoh, N. and Mita, T. Molecular cloning and expression of a novel homeobox gene AHox1 of the ascidian, <i>Halocynthia roretzi</i>	821	Conlon, F. L., Barth, K. S. and Robertson, E. J. A novel retrovirally induced embryonic lethal mutation in the mouse: assessment of the developmental fate of embryonic stem cells homozygous for the 413.d proviral integration	969
Angres, B., Müller, A. H. J., Kellermann, J. and Hausen, P. Differential expression of two cadherins in <i>Xenopus laevis</i>	829	Alberga, A., Boulay, J.-L., Kempe, E., Dennefeld, C. and Haenlin, M. The <i>snail</i> gene required for mesoderm formation in <i>Drosophila</i> is expressed dynamically in derivatives of all three germ layers	983
Danilchik, M. V. and Denegre, J. M. Deep cytoplasmic rearrangements during early development in <i>Xenopus laevis</i>	845	Torrence, S. A. Positional cues governing cell migration in leech neurogenesis	993
Serbedzija, G. N., Burgan, S., Fraser, S. E. and Bronner-Fraser, M. Vital dye labelling demonstrates a sacral neural crest contribution to the enteric nervous system of chick and mouse embryos	857	Rowe, A., Richman, J. M. and Brickell, P. M. Retinoic acid treatment alters the distribution of retinoic acid receptor- β transcripts in the embryonic chick face	1007
Noden, D. M. Origins and patterning of avian outflow tract endocardium	867	Palmer, S. J. and Burgoyne, P. S. XY follicle cells in the ovaries of XO/XY and XO/XY/YYY mosaic mice	1017
Kawabe, T. T., Rea, T. J., Flenniken, A. M., Williams, B. R. G., Groppi, V. E. and Buhl, A. E. Localization of TIMP in cycling mouse hair	877	Hashimoto, C., Gerttula, S. and Anderson, K. V. Plasma membrane localization of the <i>Toll</i> protein in the syncytial <i>Drosophila</i> embryo: importance of transmembrane signaling for dorsal-ventral pattern formation	1021
Seaver, E. C., Karlstrom, R. O. and Bastiani, M. J. The restricted spatial and temporal expression of a nervous-system-specific antigen involved in axon outgrowth during development of the grasshopper	881	Peifer, M., Rauskolb, C., Williams, M., Riggleman, B. and Wieschaus, E. The segment polarity gene <i>armadillo</i> interacts with the <i>wingless</i> signaling pathway in both embryonic and adult pattern formation	1029
Frank, E. and Sanes, J. R. Lineage of neurons and glia in chick dorsal root ganglia: analysis <i>in vivo</i> with a recombinant retrovirus	895	Christian, J. L., McMahon, J. A., McMahon, A. P. and Moon, R. T. <i>Xwnt-8</i> , a <i>Xenopus Wnt-1/int-1</i> -related gene responsive to mesoderm-inducing growth factors, may play a role in ventral mesodermal patterning during embryogenesis	1045
Hatanaka, K. and Okada, M. Retarded nuclear migration in <i>Drosophila</i> embryos with aberrant F-actin reorganization caused by maternal mutations and by cytochalasin treatment	909	Harvey, M. B. and Kaye, P. L. IGF-2 receptors are first expressed at the 2-cell stage of mouse development	1057
Chisholm, A. Control of cell fate in the tail region of <i>C. elegans</i> by the gene <i>egl-5</i>	921	Hardy, R. and Reynolds, R. Proliferation and differentiation potential of rat forebrain oligodendroglial progenitors both <i>in vitro</i> and <i>in vivo</i>	1061
Nishiyama, A., Dahlin, K. J. and Stallcup, W. B. The expression of NG2 proteoglycan in the developing rat limb	933	Glick, A. B., McCune, B. K., Abdulkarem, N., Flanders, K. C., Lumadue, J. A., Smith, J. M. and Sporn, M. B. Complex regulation of TGF β expression by retinoic acid in the vitamin A-deficient rat	1081
Cheah, K. S. E., Lau, E. T., Au, P. K. C. and Tam, P. P. L. Expression of the mouse $\alpha 1(\text{II})$ collagen gene is not restricted to cartilage during development	945	Huang, F., Dambly-Chaudière, C. and Ghysen, A. The emergence of sense organs in the wing disc of <i>Drosophila</i>	1087
Wood, A., Ashhurst, D. E., Corbett, A. and Thorogood, P. The transient expression of type II collagen at tissue interfaces during mammalian craniofacial development	955		

Ott, M.-O., Bober, E., Lyons, G., Arnold, H. and Buckingham, M. Early expression of the myogenic regulatory gene, <i>myf-5</i> , in precursor cells of skeletal muscle in the mouse embryo	1097	Jones, R. Interaction of zona pellucida glycoproteins, sulphated carbohydrates and synthetic polymers with proacrosin, the putative egg-binding protein from mammalian spermatozoa	1155
Pravtcheva, D. D., Adra, C. N. and Ruddle, F. H. Timing of paternal <i>Pgk-1</i> expression in embryos of transgenic mice	1109	Urch, U. A. and Patel, H. The interaction of boar sperm proacrosin with its natural substrate, the zona pellucida, and with polysulfated polysaccharides	1165
Parkhurst, S. M. and Ish-Horowicz, D. Mis-regulating segmentation gene expression in <i>Drosophila</i>	1121	Weeks, D. L., Walder, J. A. and Dagle, J. M. Cyclin B mRNA depletion only transiently inhibits the <i>Xenopus</i> embryonic cell cycle	1173
Hunt, P. A. Survival of XO mouse fetuses: effect of parental origin of the X chromosome or uterine environment?	1137	Su, M.-W., Suzuki, H. R., Solursh, M. and Ramirez, F. Progressively restricted expression of a new homeobox-containing gene during <i>Xenopus laevis</i> embryogenesis	1179
Schnürch, H. and Risau, W. Differentiating and mature neurons express the acidic fibroblast growth factor gene during chick neural development	1143	Index of Authors and Titles	1189
		Subject Index	1195